

Collective hormonal profiles predict group performance

Modupe Akinola^{a,1}, Elizabeth Page-Gould^b, Pranjal H. Mehta^c, and Jackson G. Lu^a

^aColumbia Business School, Columbia University, New York, NY 10027; ^bDepartment of Psychology, University of Toronto, Toronto, Canada ON M5S; and ^cDepartment of Psychology, University of Oregon, Eugene, OR 97403

Edited by Susan T. Fiske, Princeton University, Princeton, NJ, and approved July 5, 2016 (received for review March 1, 2016)

Prior research has shown that an individual's hormonal profile can influence the individual's social standing within a group. We introduce a different construct—a collective hormonal profile—which describes a group's hormonal make-up. We test whether a group's collective hormonal profile is related to its performance. Analysis of 370 individuals randomly assigned to work in 74 groups of three to six individuals revealed that group-level concentrations of testosterone and cortisol interact to predict a group's standing across groups. Groups with a collective hormonal profile characterized by high testosterone and low cortisol exhibited the highest performance. These collective hormonal level results remained reliable when controlling for personality traits and group-level variability in hormones. These findings support the hypothesis that groups with a biological propensity toward status pursuit (high testosterone) coupled with reduced stress-axis activity (low cortisol) engage in profit-maximizing decision-making. The current work extends the dual-hormone hypothesis to the collective level and provides a neurobiological perspective on the factors that determine who rises to the top across, not just within, social hierarchies.

testosterone | cortisol | groups | status | performance

Humans have always worked collaboratively in small groups to accomplish goals. In modern organizations, group work is increasing because of the growing competitive and technological demands placed on organizations and the belief that a greater variety of perspectives and approaches will lead to more innovative solutions (1–3). As a result, decades of research have focused on understanding the factors that can enhance group performance and allow some groups to outperform others. Not surprisingly, groups with structures, processes, and norms that facilitate collaboration and coordination tend to perform well (4, 5). Additionally, group composition can influence performance; groups composed of members that vary in diversity, skills, and intelligence outperform others when the appropriate structures, processes, and norms are in place (3, 6–10).

Past research on the effects of group composition on performance has focused primarily on the demographic and psychological characteristics of group members without taking into consideration the biological make-up of groups. This omission is particularly surprising because there is evidence that hormonal profiles can influence performance among individuals within a group. Indeed, past research has found that a hormonal profile characterized by high testosterone and low cortisol predicts an individual's performance and status attainment (11). Just as individuals have a hormonal profile that affects their individual performance and social status, we suggest that groups likewise can be characterized by a particular hormonal profile—their collective hormonal profile—which will influence group performance. We define a collective hormonal profile as the hormonal make-up of a group of individuals. Here, we test the hypothesis that groups with a collective hormonal profile characterized by high testosterone and low cortisol outperform and attain higher social standing among a collection of groups.

Across humans and various animal species, elevated testosterone, a steroid hormone released as the end product of the

hypothalamic–pituitary–gonadal (HPG) axis, is positively related to social dominance and behaviors that foster the achievement or maintenance of high status (12, 13). Individuals high in testosterone tend to outcompete others (14) and rise to the top of their social hierarchies, particularly in studies of nonhuman animals (15, 16). Recently, a more nuanced perspective on the link between testosterone and status attainment in humans has begun to emerge. There is growing evidence that the HPG axis acts in concert with the hypothalamic–pituitary–adrenal (HPA) stress axis to regulate status attainment within groups (11, 17). The HPA axis secretes the steroid hormone cortisol in response to stress and anxiety; thus interactions between the HPG and HPA axes can be observed by measuring testosterone and cortisol simultaneously. Specifically, the dual-hormone hypothesis posits that testosterone has a positive impact on status-attaining behavior only among individuals with low levels of cortisol (18). This hypothesis is informed by evidence that high levels of cortisol can suppress the neurobiological pathway through which testosterone impacts behavior; thus high levels of cortisol may inhibit testosterone's positive effect on dominance and status attainment (18).

The hypothesis that testosterone and cortisol interact to predict status-related behaviors for individuals has received strong empirical support in numerous contexts. For instance, individuals high in testosterone and low in cortisol have a greater number of subordinates, a measure of attained status (17); are perceived as more dominant leaders (18); are more respected by their peers (19); are more popular within their social networks (20); and engage in more competitive behavior as well as risk-taking (18, 21), a behavioral strategy associated with the attainment of social status (22, 23).

The current work explores whether the hormonal profile of a group predicts its standing among groups. We test whether the hormonal profile of high testosterone and low cortisol at the

Significance

Past research has focused primarily on demographic and psychological characteristics of group members without taking into consideration the biological make-up of groups. Here we introduce a different construct—a group's collective hormonal profile—and find that a group's biological profile predicts its standing across groups and that the particular profile supports a dual-hormone hypothesis. Groups with a collective hormonal profile characterized by high testosterone and low cortisol exhibit the highest performance. The current work provides a neurobiological perspective on factors determining group behavior and performance that are ripe for further exploration.

Author contributions: M.A. designed research; M.A. performed research; E.P.-G. and J.G.L. analyzed data; and M.A., E.P.-G., P.H.M., and J.G.L. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. Email: makinola@columbia.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1603443113/-DCSupplemental.




Fig. 1. Group performance as a function of group-level testosterone and group-level cortisol.

group performance ($b = -2.68$, $SE = 1.18$, $r = -0.27$, $P = 0.03$) (Fig. 1; see Table 2, model 2 for full statistics). When group cortisol was relatively low, group testosterone was significantly predictive of performance ($b = 1.93$, $SE = 0.93$, $r = 0.25$, $P = 0.04$) (dashed line in Fig. 1). However, group testosterone was unrelated to performance when group cortisol was high ($b = -0.51$, $SE = 1.06$, $r = -0.06$, $P = 0.64$) (solid line in Fig. 1). A robustness check excluding the covariates in the model indicated that the interaction between group-level testosterone and cortisol remained significant ($b = -3.35$, $SE = 1.42$, $r = -0.27$, $P = 0.02$) (Table 2, model 1). The positive simple slope between group testosterone and performance among groups collectively low in cortisol also remained significant ($b = 2.64$, $SE = 1.14$, $r = 0.27$, $P = 0.02$). Furthermore, group testosterone remained unrelated to performance when group cortisol was high ($b = -0.33$, $SE = 1.09$, $r = -0.04$, $P = 0.76$). The interaction between group-level testosterone and cortisol was seen not only in our aggregated measure of group performance but also in the component metrics of group performance, with the strongest of these metrics being profitability and group ranking: profitability ($b = -1027247.81$, $SE = 403289.37$, $r = -0.30$, $P = 0.01$); contracts ($b = -1.26$, $SE = 0.82$,

$r = -0.18$, $P = 0.13$); reorders ($b = -39.32$, $SE = 28.85$, $r = -0.16$, $P = 0.18$); and group ranking ($b = 9.01$, $SE = 3.57$, $r = 0.29$, $P = 0.01$). In other words, all component measures of group performance showed a trend in which group performance was highest when collective testosterone was high and collective cortisol was low. These findings demonstrate that the HPA and HPG axes play a dual role across groups, just as they do within groups.

Next we tested whether this dual-hormone interaction still predicted group performance when controlling for two personality characteristics conceptually linked to testosterone and cortisol—trait dominance and trait anxiety, respectively. An effect of collective hormonal profiles on performance in this analysis would rule out the alternative explanation that collective hormonal profiles simply reflect self-reported dispositional factors. Both dominance and anxiety were assessed via self-report several weeks before the group performance task, and unbiased group-level means of both personality traits were calculated. Neither dominance nor anxiety was correlated respectively with testosterone or cortisol (Table 1), as is consistent with prior research highlighting the relative independence of responses across nonbiological and biological systems (24–26). Furthermore, when we added group-level trait dominance, trait anxiety, and their interaction to our main regression model, the interaction between group testosterone and group cortisol remained a significant predictor of group performance ($b = -2.38$, $SE = 0.92$, $r = -0.31$, $P = 0.01$) (Table 3, model 1). This robust effect of the collective hormonal profile on group performance, when controlling for self-reported personality traits, is aligned with research suggesting that the motives captured by hormonal measures may not be easily detected by self-report measures that are meant to assess typical trait-like behavior and can apply to a variety of contexts (27). In contrast, hormonal measures may be more situational and linked to behavior in very specific contexts, which in the case of the present research are contexts that evoke status attainment. Our findings also align with theories proposing that the HPG and HPA axes influence motivation and behaviors primarily outside conscious awareness (28, 29).

The analyses thus far reveal that group-level concentrations of testosterone and cortisol predicted group performance. However, past research has found that groups with low within-group variability in a marker of prenatal androgen exposure (second-to-fourth digit ratio), which is conceptually related to social dominance and status, are less productive because of heightened intragroup conflict (30). This finding is consistent with research demonstrating that the presence of too many high-status individuals on a team can have a negative effect on group performance

Table 3. Multilevel models including trait anxiety, trait dominance, and cortisol and testosterone SDs as predictors of group performance

| Variable | Model 1 | | | | | Model 2 | | | | |
|-------------------------------|---------|------|----|-------|------|---------|------|----|-------|------|
| | Slope | SE | df | t | P | Slope | SE | df | t | P |
| Intercept | 0.18 | 0.10 | 64 | 1.76 | 0.08 | 0.19 | 0.11 | 64 | 1.68 | 0.10 |
| Cortisol, mean | 0.50 | 0.32 | 64 | 1.56 | 0.12 | 0.02 | 0.39 | 64 | 0.06 | 0.95 |
| Testosterone, mean | 0.00 | 0.66 | 64 | 0.00 | 1.00 | 0.92 | 1.09 | 64 | 0.84 | 0.40 |
| Cortisol × testosterone, mean | -2.38 | 0.92 | 64 | -2.59 | 0.01 | -2.84 | 1.17 | 64 | -2.44 | 0.02 |
| Anxiety, mean | 0.31 | 0.33 | 64 | 0.95 | 0.35 | | | | | |
| Dominance, mean | -0.46 | 0.33 | 64 | -1.42 | 0.16 | | | | | |
| Anxiety × dominance, mean | 0.44 | 1.01 | 64 | 0.44 | 0.66 | | | | | |
| Cortisol, SD | | | | | | -0.15 | 0.52 | 64 | -0.29 | 0.77 |
| Testosterone, SD | | | | | | 0.48 | 0.55 | 64 | 0.88 | 0.38 |
| Cortisol × testosterone, SD | | | | | | -0.75 | 1.67 | 64 | -0.45 | 0.65 |
| Time of day | 0.08 | 0.06 | 64 | 1.26 | 0.21 | 0.05 | 0.05 | 64 | 0.89 | 0.38 |
| No. of females | -0.19 | 0.28 | 64 | -0.68 | 0.50 | -0.03 | 0.28 | 64 | -0.11 | 0.91 |
| Group size | 0.22 | 0.15 | 64 | 1.42 | 0.16 | 0.08 | 0.15 | 64 | 0.51 | 0.61 |

$n = 74$ groups.

(31–34). To rule out the possibility that within-group hormonal variability was a key driver of our performance effects, we calculated each group's SD in testosterone and cortisol and tested whether within-group variability in testosterone, cortisol, or their interaction was related to group performance. When included in the main regression model, there were no significant effects of within-group variability in either hormone or their interaction on group performance. Further, the interaction between mean levels of testosterone and cortisol on group performance remained significant when within-group hormone levels were included in the model ($b = -2.84$, $SE = 1.17$, $r = -0.29$, $P = 0.02$) (Table 3, model 2). We also conducted exploratory analysis to test whether testosterone and cortisol unbiased means and SDs interacted to predict group performance and found no significant effects (*Supporting Information*). Additionally, we examined each group's concentration of individuals with the high-testosterone and low-cortisol profile to test whether greater concentrations of this collective hormonal profile predicted performance. We found no reliable effects of group concentration on group performance (*Supporting Information*), suggesting that the relationship between collective hormonal profiles and group performance is an emergent process that is not simply a reflection of individuals' hormonal profiles (Table S2).

The results demonstrate that groups collectively high in testosterone and low in cortisol outperform others, attaining higher standing. This finding is consistent with the dual-hormone hypothesis: Groups collectively high in testosterone may have a heightened status drive, but this status drive results in optimal decision-making only when coupled with low cortisol, a potential indicator of reduced stress and attenuated behavioral inhibition. Indeed, recent evidence suggests that increased testosterone combined with reduced cortisol may elevate an individual's status by promoting rational decision-making focused on maximizing financial profits (35). Our findings extend this research to the group level and suggest that groups with a propensity for status attainment, but without inhibition, engage in profit-maximizing decision-making strategies.

Our findings offer several avenues for future research. For instance, future studies should try to capture the emergent process through which an individual's neurobiology is transformed in an adaptively synergistic way to affect group performance. It is also critical to consider other neurobiological profiles that might influence group performance. Because anabolic hormones, such as dehydroepiandrosterone (DHEA), can counterregulate catabolic hormones (e.g., cortisol) and are indicative of an adaptive, thriving response during acute stress (36), we may expect to find that the interaction of group-level DHEA and cortisol predicts group performance under acute stress. This example, combined with our findings, highlights the importance of examining both the linear and interactive effects of collective hormones on group outcomes to capture the context-dependent nature of hormones (18), particularly as they relate to HPA and HPG axis activation. Additionally, although here we focus on resting hormone levels, examining hormonal reactivity and hormonal covariation among group members is another important direction for future research.

Our theorizing focuses on status attainment motivation as the key mechanism underlying our group performance effect; however, future studies should examine group-process measures to advance our knowledge of how status attainment motivations manifest themselves in group dynamics. There is evidence that the combination of elevated testosterone and reduced cortisol produces greater social rapport in interdependent decision-making tasks (35). This past finding suggests that groups in the present study with a collective hormonal profile of high testosterone and low cortisol may have experienced greater intragroup cohesion. Because intragroup cohesion is a critical determinant of group performance (37–40), we suspect that high testosterone/

low cortisol groups produced more favorable performance outcomes partly because they were more cohesive.

The current research goes beyond past work that has focused on observable behaviors of the group or on psychological characteristics of individuals within the group as predictors of group performance without paying attention to biological factors at the collective level. By showing that groups collectively high in testosterone and low in cortisol outperform other groups, we provide a broader perspective on the processes at play in determining which groups rise to the top across and not just within social hierarchies. In doing so, we both establish evidence in support of the dual-hormone hypothesis at the collective level and open up additional questions regarding the neurobiology of group behavior and performance.

Materials and Methods

Participants. Participants were 370 Master in Business Administration (MBA) students (63.8% male; 36.2% female) enrolled in an introductory organizational behavior course and an operations management course at a business school in the northeastern United States. The average age of the sample was 27.44 y ($SD = 1.93$ y); 157 participants (42.4%) were born outside the United States. The ethnic composition of our sample was diverse: 54.9% white, 16.5% Asian, 10.8% Hispanic, 9.4% South Asian, 4.6% black, 1.4% Southeast Asian, and 2.4% other. Participants provided informed consent, and the Institutional Review Board of Columbia University approved all materials and procedures in our study.

Procedure. Participants were randomly assigned to 74 groups ranging in size from three to six people. These groups met regularly during the semester (6 wk) and were required to work on a variety of tasks together outside of class. There were no formally assigned positions or roles within the groups, allowing groups to develop roles and form status hierarchies naturally over time (41). Additionally, there was little variance across groups in gender composition, because the MBA sample was predominantly male. Specifically, 1.4% of the groups had no women, 18.9% of the groups had one woman, and the remaining 79.7% had two women. Furthermore, gender composition did not moderate our observed effects [$b = 0.15$, $SE = 4.47$, $t(64) = 0.03$, $r < 0.01$, $P = 0.85$]. The lower-level interaction between mean levels of testosterone and cortisol was still a significant predictor of group performance when gender was considered as a moderator instead of as a covariate [$b = -2.74$, $SE = 1.28$, $t(64) = -2.14$, $r = -0.26$, $P = 0.04$].

At the start of the semester, during their organizational behavior course, participants completed a survey assessing trait dominance. Two weeks later, they provided a saliva sample and completed a survey assessing trait anxiety. One week after providing the saliva sample, participants engaged in a competitive group decision-making task in their operations management course in which their performance was measured relative to other groups in the class at two time points.

Measures.

Testosterone and cortisol. Participants provided one saliva sample at the start of class during their assigned class time of 9:00 AM (early morning session), 10:45 AM (late morning session), 2:15 PM (early afternoon session), or 4:00 PM (late afternoon session). Participants first completed informed consent and biological questionnaires and then, after a 15-min rest period, provided a saliva sample that was later assayed for testosterone and cortisol. Saliva samples were frozen until the study was completed and then were shipped overnight on dry ice to a laboratory in College Park, PA. Saliva samples were assayed for testosterone and cortisol using a highly sensitive enzyme immunoassay (Salimetrics). Average intra- and interassay coefficients of variation were 2.5% and 5.6%, respectively, for testosterone and 3.5% and 5.1%, respectively, for cortisol. Before analysis, both hormone measures were log-transformed to account for significant positive skew, and mean levels were centered around their grand means (see *Supporting Information* and Table S3 for raw hormone values). Finally, group-level testosterone and cortisol profiles were calculated from the best linear unbiased predictors of testosterone and cortisol levels within each group (see *Data Analysis Strategy*), because groups varied in size from three to six members. We also calculated group-level SDs in testosterone and cortisol for each group.

Group performance. The group decision-making task was a computerized exercise (Littlefield Labs, Responsive Learning Technologies) simulating the supply-chain process of a series of laboratories providing blood testing to its

34. Tiedens LZ, Fragale AR (2003) Power moves: Complementarity in dominant and submissive nonverbal behavior. *J Pers Soc Psychol* 84(3):558–568.
35. Mehta PH, Mor S, Yap AJ, Prasad S (2015) Dual-hormone changes are related to bargaining performance. *Psychol Sci* 26(6):866–876.
36. Epel ES, McEwen BS, Ickovics JR (1998) Embodying psychological thriving: Physical thriving in response to stress. *J Soc Issues* 54(2):301–322.
37. Beal DJ, Cohen RR, Burke MJ, McLendon CL (2003) Cohesion and performance in groups: A meta-analytic clarification of construct relations. *J Appl Psychol* 88(6):989–1004.
38. Evans CR, Dion KL (1991) Group cohesion and performance a meta-analysis. *Small Group Res* 22(2):175–186.
39. Gully SM, Devine DJ, Whitney DJ (1995) A meta-analysis of cohesion and performance effects of level of analysis and task interdependence. *Small Group Res* 26(4):497–520.
40. Mullen B, Copper C (1994) The relation between group cohesiveness and performance: An integration. *Psychol Bull* 115(2):210–227.
41. Moreland RL, Levine JM (1988) Group dynamics over time: Development and socialization in small groups. *The Social Psychology of Time: New Perspectives*, eds McGrath J, Joseph E (Sage, Thousand Oaks, CA), pp 151–181.
42. Wiggins JS, Trapnell P, Phillips N (1988) Psychometric and geometric characteristics of the Revised Interpersonal Adjective Scales (IAS-R). *Multivariate Behav Res* 23(4): 517–530.
43. Anderson C, Berdahl JL (2002) The experience of power: Examining the effects of power on approach and inhibition tendencies. *J Pers Soc Psychol* 83(6):1362–1377.
44. Watson D, Clark LA, Tellegen A (1988) Development and validation of brief measures of positive and negative affect: The PANAS scales. *J Pers Soc Psychol* 54(6): 1063–1070.
45. Mehta PH, Josephs RA (2011) Social endocrinology: Hormones and social motivation. *The Handbook of Social Motivation*, ed Dunning D (Psychology, New York).
46. Croon MA, van Veldhoven MJPM (2007) Predicting group-level outcome variables from variables measured at the individual level: A latent variable multilevel model. *Psychol Methods* 12(1):45–57.
47. R Core Team (2015) *R: A Language and Environment for Statistical Computing* (R Found Stat Comput, Vienna).
48. Punj G, Stewart DW (1983) Cluster analysis in marketing research: Review and suggestions for application. *J Mark Res* 20:134–148.